

THE EVOLUTION OF THE BL LAC OBJECTS

A. CAVALIERE¹ AND D. MALQUORI^{2,3}

Accepted for publication in ApJ Letters

ABSTRACT

The BL Lac population is set apart from the rest of the active galactic nuclei by a number of peculiar features, including remarkably little signs of cosmological evolution. We focus on this feature, and take hint from the observations to reconsider extraction of the primary energy from a Kerr hole via the Blandford-Znajek (BZ) process; in fact, we stress how this can power the emitting jets of the BL Lac objects in a regime of low accretion. We show that the timescale for the BZ process to exhaust the rotational energy of the hole spans many gigayears in all such sources. We compute scales long and uniform enough to yield an intrinsically slow population evolution, giving rise to number counts of “Euclidean” or flatter shape; even longer timescales apply to the additional, analogous process involving the inner accretion disk directly. Such slow evolution takes place in ambients and at epochs ($z \lesssim 1$) where interactions of the host galaxies with companions become rare or weak. On the other hand, at higher z , we argue that frequent interactions of the hosts within groups enhanced, each for some 10^{-1} Gyr, the accretion rates and the emissions from the disks, attaining power levels that are typical of the radio-loud quasars. But such interactions were also likely to disorder the large-scale magnetic field in the disk and to decrease the angular momentum of the hole, thus limiting the lifetime of all BZ emissions to under a few gigayears. Then the stronger evolution of the flat-spectrum radio-loud quasars was driven, similar to the radio-quiet ones, by the interaction rate decreasing rapidly with z .

Subject headings: BL Lacertae objects: general – galaxies: active – galaxies: evolution – radiation mechanisms: nonthermal

1. INTRODUCTION

The BL Lacertae objects among the active galactic nuclei (AGNs) exhibit a number of peculiarities (see Urry & Padovani 1995, Sambruna 1999, and references therein), namely: (a) lack of emission lines with equivalent width exceeding 5 Å; (b) a pure nonthermal continuum; (c) flat-spectrum radio emission, and powerful γ -rays that extend up to TeV energies in some objects; (d) rapid variability; and (e) strong and highly variable optical polarization. To some extent, they share the three latter features with the flat-spectrum radio-loud quasars (FSRQs).

The current understanding links the peculiarities *c*, *d*, and *e* to synchrotron and inverse Compton emissions from a relativistic jet with bulk Lorentz factors $\Gamma \sim 3 - 10$, viewed at a small angle of order Γ^{-1} (see Ghisellini et al. 1993). On the other hand, the peculiarities *a* and *b* are widely related to a dearth of gas surrounding the activity center at both parsec and 10 μ pc scales, and/or to the lack of an adequate nonbeamed continuum. An additional peculiarity is constituted by the behavior of the population, which shows *little or no* signs of evolution (see Stickel et al. 1991; Wolter et al. 1991; Bade et al. 1998), at striking variance with the rest of the AGNs.

To us, the peculiarities *a* and *b* suggest that current accretion is *low* in BL Lac objects, providing little gas and a weak ionizing continuum. Taking up previous proposals (see Blandford 1993 and references therein), we consider the main power source for the BL Lac emissions to be provided by the extraction in electromagnetic form of *rotational* energy, stockpiled in the central black hole and

associated with the inner accretion disk. We develop this view to explain the other BL Lac peculiarities, and specifically the weak evolution.

2. EVOLUTIONARY BEHAVIOR

Once Doppler enhancements of order Γ^{3-4} are accounted for, the observations of BL Lac objects (see Sambruna, Maraschi & Urry 1996), indicate typical rest frame luminosities $L \lesssim 10^{44}$ ergs s⁻¹, which are moderate on the quasar standards; meanwhile, the rotational energies extractable from Kerr holes (KHs) are around $E \sim 5 \cdot 10^{61} M_8$ ergs. So their exhaustion would take in individual objects times $\tau \sim E/L$ of many gigayears, which are quite *longer* than the flashes lasting about 10^{-1} Gyr of standard quasars fueled by accretion (see Cavaliere & Vittorini 1998). In the few instances of larger L , the Doppler factors inferred from the data in the framework of the synchrotron self-Compton radiation – which constitute lower bounds anyway – are particularly low and uncertain pending confirmation from superluminal velocities (P. Padovani 1998, private communication). The long debated kinetic component of the jets is unlikely to increase the power budget of the typical BL Lac objects beyond some 10^{45} ergs s⁻¹ (Ghisellini & Celotti 1998).

The population-averaged $\langle L(t) \rangle$ is constituted by the convolution of the individual behaviors. When these are all similar, as we shall show below, $\langle L(t) \rangle$ also shares the same timescale τ . Heuristically, a long τ will imply little population evolution and, therefore, relatively flat number counts, or equivalently V/V_{max} close to 0.5.

¹ Astrofisica, Dipartimento di Fisica, 2a Università degli studi di Roma, Roma, 00133, Italy

²Dipartimento di Astronomia, Università di Padova, Padova, 35122, Italy

³Institut d’Astrophysique de Paris (CNRS), Paris, 75014, France

A quantitative statement can be made in simple terms on recalling that the integral counts $N(> S)$ at high fluxes S may be cast into the form (see Weinberg 1972):

$$N(> S) \propto S^{-3/2} [1 - C(S_o/S)^{1/2} + 0(S^{-1})] . \quad (1)$$

In turn, C can be expressed (Cavaliere & Maccacaro 1990) in terms of the time scale τ_e for population changes by

$$C = \frac{3 D_o < l^2 >}{4 R_H < l^{3/2} >} [2(1 + \alpha) - \frac{1}{H_o \tau_e}] . \quad (2)$$

Here $R_H = 3000 h^{-1}$ Mpc is the Hubble radius with the Hubble constant $h = H_o/100$ km s $^{-1}$ Mpc $^{-1}$, $D_o \equiv (P_o/4\pi S_o)^{1/2} \approx 200 h^{-1}$ Mpc in terms of the power and flux of the brightest BL Lac objects, $< l^n >$ is the (normalized) n -th moment of the luminosity function (LF), and α is the spectral index.




FIG. 1 – The runs of $A(t)$ and of $L(t)$ for $\dot{M} = \text{const}$, corresponding to initial normalized accretion rates $\dot{m}_o = 5 \cdot 10^{-2}$ (upper curves), 10^{-2} (middle curves), and 10^{-3} (lower curves). Note the convergent behaviors at the lowest rates.

The evolutionary rate $\tau_e^{-1} = \tau_D^{-1} + (\beta - 1)\tau_L^{-1}$ comprises two contributions that are due to density and to luminosity evolutions, respectively; the latter is amplified by a steep LF (modeled with $L^{-\beta}$ in the relevant range), and τ_D is negative in case of negative evolution. From eq. (2) it is seen that the bright counts tend to the “Euclidean” shape $N(> S) \propto S^{-3/2}$ or flatter even when $\beta \gtrsim 2.5$, and even in the presence of positive evolution, provided this is slow enough to satisfy $\tau_e H_o \geq [2(1 + \alpha)]^{-1}$.

3. A SIMPLE MODEL: BEAMS FROM KERR HOLES

We begin with considering the specific model where the primary power is extracted from the rotational energy of a KH, following Blandford & Znajek (1977). They tackled a stationary, axisymmetric and force-free magnetosphere surrounding the KH, and derived the scaling

$L_h \propto B^2 r_h^2 A^2$ for the power extracted from the KH *itself* in the form of Poynting flux; here $A \leq 1$ denotes the angular momentum of the hole, J , in units of GM^2/c , and B is the magnetic field threading the horizon at $r_h = [1 + (1 - A^2))^{1/2}] GM/c^2$.

The strength of the field that is sustainable at the horizon constitutes a delicate issue. A recent reappraisal by Ghosh & Abramowicz (1997) has been based on continuity of the magnetic pressure at the inner edge of the accretion disk and on numerical computations – which are self-consistent, if with dynamic range limited to small and medium scales – of its MHD and thermal structure. They find that $B^2/8\pi \sim \alpha p$ holds for standard α -disks in terms of the maximal pressure at the inner rim; they (conservatively) evaluate the resulting power to read

$$L_h \simeq \frac{2 \cdot 10^{44} M_8 A^2 \text{ ergs s}^{-1}}{8 \cdot 10^{42} M_8^{11/10} \dot{m}_{-4}^{4/5} A^2 \text{ ergs s}^{-1}} , \quad (3)$$

when the inner region is dominated, respectively, by radiation [$p \propto (\alpha M)^{-1}$] or by gas pressure [$p \propto (\alpha M)^{-9/10} \dot{m}^{4/5}$]. We recall from Novikov & Thorne (1973) that only when the accretion rate in Eddington units exceeds $\dot{m} \equiv \dot{M} c^2 / L_E \simeq 3 \cdot 10^{-4} M_8^{-1/8}$ (for $A = 1$) does the transitional radius $r_* \propto \dot{M}^{16/21}$ (separating the radiation-dominated region from the gas-dominated region) exceed the radius r_{ms} of the marginally stable orbit so that an inner, radiation pressure dominated region in fact occurs. Then the ratio of the Blandford-Znajek (BZ) emission to the thermal luminosity scales as $L_h/L_{th} \approx 3 \cdot 10^{-2} A^2 / \dot{m}$. Thus, in the range $10^{-3} \lesssim \dot{m} \lesssim 10^{-2}$, the rest frame power L_h is adequate to sustain the nonthermal, beamed emissions of BL Lac objects and to match or dominate the thermal component L_{th} .

As for the evolutionary behavior of the population, we first consider the time dependence of $L_h(t)$ for an individual source; this is related, by equations (3), to the behaviors of $M(t)$ and of $J(t)$. The key point is that accretion of J also takes place along with $\dot{M}$, and is always important, as first pointed out by Bardeen (1970), being related to $\dot{M}$ by the conditions in the disk at r_{ms} .

In fact, the run of $M(t)$ is described by

$$\frac{dM}{dt} = e_{ms} \dot{M} - L_h / c^2 , \quad (4)$$

where $e_{ms} \simeq 1$ is the adimensional specific energy at r_{ms} ; actually, $e_{ms} = 0.58$ holds for $A = 1$, rising to 0.94 for $A = 0$ (de Felice 1968; Bardeen 1970). On the other hand, the run of $J(t)$ is coupled to the former by

$$\frac{dJ}{dt} = j_{ms} \dot{M} - \frac{L_h}{k \Omega_h} \quad (5)$$

(Malquori 1996; Moderski & Sikora 1996). Here j_{ms} is the specific angular momentum at r_{ms} , with values close to $2 e_{ms} c r_h$; moreover, $\Omega_h \equiv A c / 2 r_h$, while $k = 1/2$ for maximum power extraction.

On the right-hand side of equation (5) the accretion of J can be balanced out by the loss due to the BZ process, causing dJ/dt to vanish for values of A around $50 \dot{m}$, given $10^{-3} \lesssim \dot{m} \lesssim 10^{-2}$. Instead, $A(t) \propto J(t)/M^2(t)$ decreases monotonically (see Fig. 1). Correspondingly, $L_h(t)$

(shown in Fig. 1 with all relativistic factors and detailed A -dependencies in place) decreases steeply at first because of the relativistic factors, but then it tends to level off (or to increase again for high $\dot{M}$) because of the hole mass increase; in between, the decrease takes place on the scale $\tau \equiv L/\dot{L} \simeq J/2\dot{J}$. Such behavior and scale are *similar* for all objects at low $\dot{m}$ where the BZ power dominates.

In sum, the sources powered by KHs fade slowly, but remain above levels $L_{ho}/10 \sim 10^{44} \text{ erg s}^{-1}$ that are relevant for BL Lac objects for times $\tau \gtrsim 8 \text{ Gyr}$. So their population undergoes a slow luminosity evolution; the condition $\tau_e H_o \simeq [2(1 + \alpha)]^{-1}$ tends to hold especially for $h \simeq 0.7$, and the bright counts in *any* band ought to be quite *flatter* than with the quasars.

As an example, we compute in detail eqs. (1) and (2) for the radio band on the basis of the above luminosity evolution alone. In the radio, the BL Lac counts are arguably least sensitive to the spectral changes with power proposed by Fossati et al. (1998; see also Georgenopoulos & Marscher 1998 and Rector et al. 1999). There $\alpha \simeq 0$ and $\beta \simeq 2.5$ apply (we refer to the model LF computed by Padovani & Giommi 1995), and the condition $\tau_e H_o \simeq 1/2$ for nearly Euclidean counts is closely met. In fact, we find the shape shown in Figure 2 by the solid line; this is easily consistent with the best fit (*dashed line*) to the counts by Stickel et. al. (1991), a published sample that is complete although it is affected by much statistical variance. A flatter LF will yield counts flatter still. On the other hand, a flatter LF, but with fast evolution, would yield steep counts as illustrated by the dotted line in Figure 2.

Actually, our results provide an upper limit to the counts, since the values of τ evaluated above provide minimal timescales while an additional negative density evolution is likely to occur, as discussed next. More extensive and deeper data will be released soon (see Giommi, Menna & Padovani 1999 and Rector et al. 1999) and will provide more challenging evolution indicators for our prediction.

4. DISCUSSION AND CONCLUSIONS

All peculiarities of the BL Lac objects are consistent with the extraction in *electromagnetic* form of the *rotational* energy stockpiled in a KH, in a regime of *low* accretion. In fact, a primary Poynting flux, as produced via the BZ process by the KH itself, is attractive (Ghisellini & Celotti 1998) because it is energetically adequate and is conducive to jets with bulk $\Gamma \sim 10$; by way of the implied large-scale electric fields, it is conducive also to distributed and continuous acceleration up to isotropic electron energies in the range of 10^2 GeV . Thus, highly beamed, non-thermal (and polarized) continua extending into the TeV γ -ray range can be generated; if a jet is oriented close to our line of sight, high apparent power and fast time variability will be observed (Begelman et al. 1984). In turn, *low* current accretion rates, $\dot{m} \lesssim 10^{-2}$, are consistent with negligible dilution by thermal continua (then also unable to degrade the TeV γ -rays) and with emission lines of small equivalent width. But our thrust has been to show how this view leads us to predict intrinsically *slow* population evolution.

Actually, an additional MHD emission L_d is produced by the inner disk (Blandford & Znajek 1977; Livio, Ogilvie & Pringle 1999). This scales still with B^2 , but also with

$r_*^{3/2} r_h^{1/2}$ so that the ratio $L_d/L_h \propto (r_*/r_h)^{3/2} \propto \dot{M}^{1.15}$; the coefficient is independent of $A^2(t)$ and somewhat larger than in eq. (3a). Consideration of L_d does not affect materially our evaluations as long as $\dot{m} \lesssim 10^{-2}$, and so $L_d \sim L_h$, which is appropriate to the BL Lac objects. Then the KH and the disk are dynamically and magnetically linked nearly in one system in which – within (uncertain) factors of a few – the hole still may be regarded as the energetic flywheel.




FIG. 2 – BL Lac counts at 5 GHz, from Stickel et al. 1991. The dashed line reproduces their best fit $N(> S) \propto S^{-1.6}$ (the actual slope is 1.58 ± 0.44 at a 1σ level). The solid line gives our result for pure luminosity evolution with $\tau = 8 \text{ Gyr}$, assuming constant L_{radio}/L_h and $\beta = 2.5$. The dotted line represents for comparison a case where $\beta = 2.2$ and $\tau = 2 \text{ Gyr}$.

The stand in this context of the other Blazars, the FSRQs, is interesting as such objects exhibit features similar to the BL Lacs, less pure but with larger outputs. Tapping the rotational energy in order to produce relativistic jets, blazing beams, and nonthermal luminosities is tantalizing also for these objects, but there are also important differences.

Now the maximal KH power L_h from eq. (3a) is inadequate – with masses limited to $M \lesssim 410^9 h^{-1} M_\odot$ (see Magorrian et al. 1998) – to match powers of $10^{46} h^{-2} \text{ erg s}^{-1}$ or more; a larger disk component L_d to the BZ emission is cogently called for. In fact, $L_d \gg L_h$ now holds (with the KH now slaved to the disk) since $\dot{m} \sim 1$; we know this is to hold for the FSRQs, because their emission lines and thermal components, which dilute the BL Lac-like peculiarities to approach the standard quasar activity, do signal currently *high* accretion rates. We also expect the larger disk component in FSRQs to imply – relative to the BL Lac conditions – more plasma lifted up and electric fields more accurately screened out; then the electrons not only may loose their energy faster at such larger total outputs (Fossati et al. 1998), but are also accelerated to lower energies, which is another, intrinsic reason to emit

softer γ -rays.

Since the BZ emissions are marked by long timescales at constant $\dot{M}$, the long persistence of radio-emitting, rotationally driven jets might be seen as statistically embarrassing (see Moderski, Sikora & Lasota 1998). However, we argue that the actual timescales of the FSRQs are much shorter because of the boundary conditions set to such high values of $\dot{M}$ by the host galaxy. For the sake of this argument, we shall discuss first the scenario (most constraining to us) where the BL Lac objects are descendants of FSRQs.

We recall from Cavaliere & Vittorini (1998) that *recurrent* interactions and merging events of the E host galaxies with satellites or with group companions (for which there is plenty of morphological and kinematical evidence; see Bertola & Corsini 1998) are necessary to cause substantial losses of angular momentum, $|\Delta j|/j \sim 1$, of the gas residing in either interaction partner; such events every $1 - 2$ Gyr trigger for some 10^{-1} Gyr gas inflow toward the nucleus, which is conducive to $\dot{M}$ and also to accretion of angular momentum onto the hole. We concur with Moderski, Sikora & Lasota (1998) that the *sign* of such imported momentum relative to the previous one makes a difference as for disk structure and for the KH rotation. We stress that each strong interaction has a $1/2$ chance to accrete a component opposite to the initial $\mathbf{J}$ of the KH and of the disk, thus wrecking the latter's large-scale magnetic field and spinning the former down, effectively quenching their emissions. When the disk is rebuilt, the internal conditions for large-scale $\mathbf{B}$ and those in the host galaxy for beam propagation (see Livio 1999) may not be met again.

Thus we hold that the probability is small for combining persistent extraction of BZ power with frequent episodes of high $\dot{m}$. But the quasar evolution since $z \approx 3$ is governed just by high- $\dot{m}$ episodes driven by *interactions*; these are initially frequent and strong in the era of group formation, but then drop off in rate and strength with decreasing z as groups are reshuffled into clusters (Cavaliere & Vittorini 1998). As long as such interactions are effective, they will intermittently switch off all BZ outputs from the KH and the disk.

We conclude that strong evolution and long-lived extraction of BZ power entertain a *complementarity* relationship. Indeed, the rotational power and the beams it feeds persist safely for many gigayears only in a regime of permanently low accretion, which is what prevails for $z \lesssim 1.5$, after a "last interaction" setting $A \simeq 1$; such conditions will lead

to the BL Lac phenomenon. At higher z , the most likely activity pattern was that corresponding to FSRQs, with dominant thermal and disk BZ emissions limited to a few gigayears by initially frequent, but then subsiding, interactions; the evolution of such sources is then expected to be close (cf. Goldschmidt et al. 1999) to the standard quasars', being driven by the same z -dependent $\langle \dot{m} \rangle$ only somewhat slowed down by the beam attempts to persist in between interactions. On the other hand, if it is maintained (see Jackson & Wall 1999) that BL Lacs and FSRQs belong to different and secluded lineages, then little constraints will arise from FSRQs for the the present view of the astrophysics and evolution of BL Lac objects.

Finally, enlarging our view of this recently rejuvenated field above details and beyond the theoretical dream of a relativistic homopolar inductor, we summarize and emphasize the two key issues in this context. First and concerning the BL Lac population, up to now flat counts and $V/V_{max} \simeq 0.5$ persistently signal a *weaker* evolution than with the quasars. This indicates either a slower luminosity evolution with timescales $\tau_L/(\beta - 1)$ of several gigayears, as considered above, or a negative density evolution with $\tau_D < 0$, or some mix of the two (some negative density evolution is also implied from BL Lac birth since $z \approx 1.5$). Second, concerning the source structure, the current wisdom on disks (but ADAF or ADIOS flows, possibly setting in at low $\dot{m}$, as discussed by Blandford & Begelman 1998, have yet to be firmly fit in) has that the disk radiation pressure is independent of accretion for $\dot{m}$ above some 10^{-3} , since then $p \propto T^4$ holds in the inner region. *As long as* good coupling of radiation, particles and large-scale magnetic fields can be retained, this leads to a *large* ratio $(L_h + L_d)/L_{th} \propto 1/\dot{m}$; in addition, as $\dot{m}$ approaches 10^{-3} from above, r_* shrinks to approach r_{ms} , and *comparable* $L_h \sim L_d$ obtain. Such conditions are best suited to yield pure nonthermal radiation of power moderate but extending up to TeV γ -rays, and by the same token they yield persistent beams with intrinsically weak evolution, precisely the features marking out the BL Lac objects.

ACKNOWLEDGMENTS

We thank V. D'Elia and P. Padovani for several useful discussions, and our referee for helpful comments. A.C.'s share of the work was completed while he visited the Asiago Observatories. The grants were from ASI and MURST.

REFERENCES

- Bade, N., et al. 1998, A&A, 334, 459
 Bardeen, J. 1970, Nature 226, 64
 Begelman, M.C., Blandford, R.D. & Rees, M.J. 1984, Rev. Mod. Phys., 56, 255
 Bertola, F., & Corsini, E.M. 1998, preprint (astro-ph/9802314)
 Blandford, R.D. 1993, in Astrophysical Jets, ed. D. Bulgarella, M. Livio, & C.P. O'Dea (Cambridge: Cambridge Univ. Press), 15
 Blandford, R.D., & Begelman, M.C. 1998, MNRAS, 303, L1
 Blandford, R.D. & Znajek, R.L. 1977, MNRAS, 179, 433
 Cavaliere, A., & Maccacaro, T. 1990, ApJ, 358, 1
 Cavaliere, A., & Vittorini, V. 1998, in ASP Conf. Ser. 146, The Young Universe: Galaxy Formation and Evolution at intermediate and High Redshift, ed. S. D'Odorico, A. Fontana & E. Giallongo (San Francisco: ASP), 26
 de Felice, F. 1968, Nuovo Cimento, 57B, 351
 Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G. 1998, MNRAS, 299, 433
 Ghisellini, G., & Celotti, A. 1998, MNRAS, submitted
 Ghisellini, G., Padovani, P., Celotti, A., & Maraschi, L. 1993, ApJ, 407, 65
 Ghosh, P., & Abramowicz, M.A. 1997, MNRAS, 292, 887
 Georganopoulos, M., & Marsher, A.P. 1998, ApJ, 506, 621
 Giommi, P., Menna, M. T. & Padovani, P. 1999, in ASP Conf. Ser. 159, The BL Lacertae Phenomenon, ed. L.O. Takalo & A. Sillanpaa (San Francisco: ASP), in press
 Goldschmidt, P., Kukula, M.J., Miller, L., & Dunlop, J.S. 1999, ApJ, 511, 612
 Jackson, C.A., & Wall, J.V. 1999, MNRAS, in press (astro-ph/9808054)
 Livio, M. 1999, Phys. Rep., in press (STScI preprint No. 1261)
 Livio, M., Ogilvie, G.I., & Pringle, J.E. 1999, ApJ, 512, 100
 Magorrian, J., et al. 1998, AJ, 115, 2285
 Malquori, D. 1996, Laurea thesis, University of Padova
 Moderski, R., & Sikora, M. 1996, MNRAS, 283, 854

- Moderski, R., Sikora, M., & Lasota, J.P. 1998, MNRAS, 301, 142
Novikov, I., & Thorne, K.S. 1973, in Black Holes, ed. B. De Witts & C. De Witts (New York: Gordon & Breach), 343
Padovani, P., & Giommi, P. 1995, ApJ, 444, 567
Rector, T., Stocke, J., Perlman, E. & Gioia, I. 1999, Abstract for the AAS Meeting, 193, 107.13
Sambruna, R. 1999, in ASP Conf. Ser. 159, The BL Lacertae Phenomenon, ed. L.O. Takalo & A. Sillanpaa (San Francisco: ASP), in press (astro-ph/9811162)
Sambruna, R., Maraschi, L., & Urry, M. 1996, ApJ, 463, 444
Stickel, M., Padovani, P., Urry, M., Fried, W., & Kühr, H. 1991, ApJ, 374, 431
Urry, M., & Padovani, P. 1995, PASP, 107, 803
Weinberg, S. 1972, Gravitation and Cosmology (New York: Wiley)
Wolter, A., Gioia, I., Maccacaro, T., Morris, S., & Stocke, J. 1991, ApJ, 369, 314

E-mail: cavaliere@roma2.infn.it; malquori@pd.astro.it